



Measure Calculations

1	Proportion Measure Calculations	1
1.1	Elements of a Proportion Measure	4
1.2	Performance Rates	4
1.3	Proportion Measure Examples	4
2	Continuous Variable Calculations	7
2.1	Elements of Continuous Variable Measures	7
2.2	Continuous Variable Aggregate Calculations.....	9
2.3	CV Measure Example	9
3	Ratio Measure Calculations	10
3.1	Elements of a Ratio Measure.....	10
3.2	Ratio Measure Aggregate Calculations.....	13
3.3	Ratio Measure Example	13
4	Key Points.....	14
	References.....	16

There are multiple ways to calculate quality measures①. This document provides information about the calculations of proportion①, continuous variable (CV)①, and ratio① measures, and supplements the information found in the *Blueprint*, Chapter 5, Measure Specification.

1 PROPORTION MEASURE CALCULATIONS

This section provides guidance on the calculation and interpretation of proportion measures.

A proportion is a measure in which the numerator① is a subset (or part) of the denominator① and is written as $a/(a+b)$ (Figure 1). A proportion measure differs from a ratio measure because in a ratio measure, the numerator is not a subset of the denominator population (refer to the *Blueprint* Chapter 5.1, Measure Specifications by Measure Category). Although the numerator and denominator populations may be related or may overlap for ratio measures, these populations do not have a superset/subset relationship. Figure 2 diagrams the flow from target/initial population① to numerator and denominator.




Figure 1. Proportion Measure Populations




Figure 2. Population Criteria Definitions for Proportion Measures (Health Level Seven International® [HL7], n.d.)

1.1 ELEMENTS OF A PROPORTION MEASURE

From these relationships and definitions, the measure developer should use these sequential steps to determine whether a patient falls into a given population:

1. Target/Initial population (IP): Identify those patients who meet the IP criteria.
2. Denominator (DENOM): Identify the subset of the IP that meet the DENOM criteria.
3. Denominator exclusion① (DENEX): Identify the subset of the DENOM that meet the DENEX criteria. Remove those that are patients from the denominator as exclusions. After removal of these patients, the remaining subset reflects the group of patients evaluated for the numerator criteria.
4. Numerator (NUMER): Identify those in the DENOM and not in the DENEX that meet the NUMER criteria. In proportion measures, the numerator criteria are the processes or outcomes expected for each patient, procedure, or other unit of measurement defined in the denominator.
5. Numerator exclusion① (NUMEX): Identify the subset of the NUMER that meet the NUMEX criteria. Use NUMEX to define instances not included in the numerator data.
6. Denominator exception① (DENEXCEP): Identify those in the DENOM and not in the DENEX and NOT in the NUMER that meet the DENEXCEP criteria. Remove these patients from the IP and/or DENOM.

1.2 PERFORMANCE RATES

Specific programs may require reporting of performance rates. The performance rate is the number of patients meeting NUMER criteria (accounting for exclusions), divided by patients in the DENOM (accounting for exclusion and exception). Calculate the performance rate using the formula

$$\text{Performance rate}^1 = (\text{NUMER} - \text{NUMEX}) / (\text{DENOM} - \text{DENEX} - \text{DENEXCEP})$$

Exception: 0% Performance Rates

- Inverse measures①: For inverse measures, a lower rate indicates better performance; measure developers should consider a 0% performance rate as satisfactorily reporting (e.g., 30-day mortality rate for acute myocardial infarction).
- Null Scores①: If the measure is not applicable for all patients within the sample, the performance rate would be 0/0 (null) and is considered satisfactorily reporting or having data completeness. Do not count performance exclusion Quality Data Codes (QDCs) in the performance denominator.

1.3 PROPORTION MEASURE EXAMPLES

This section provides examples of the mathematical relationships between populations in proportion measures.

Example #1: Proportion Measure

¹ Also referred to as the raw rate.

A fictitious proportion measure defines the population criteria.

- IP: all patients aged 65 years and older with an active diagnosis of diabetes mellitus
- DENOM: equals IP
- DENEX: bilateral blindness
- NUMER: dilated eye exam for diabetic retinopathy
- NUMEX: none
- DENEXCEP: bed confinement status in a community where mobile eye exam imaging is unavailable

Mr. Jones is 75 years old and has an active diagnosis of diabetes. There is no mention of blindness in his chart. He has a documented dilated eye exam for diabetic retinopathy.

- (IP = YES) Mr. Jones meets the IP criteria.
- (DENOM = YES) Mr. Jones meets the DENOM criteria.
- (DENEX = NO) By the positive evidence principle, Mr. Jones does not meet the DENEX criteria.
- (NUMER = YES) Mr. Jones meets the NUMER criteria.
- (NUMEX = NO)
- (DENEXCEP = NO) no evaluation of the DENEXCEP because the patient met the NUMER criteria.

Mr. Smith is 75 years old and has an active diagnosis of diabetes. There is no mention of blindness in his chart. There is no mention of a dilated eye exam in his chart. There is no mention in his chart that he is bed bound.

- (IP = YES) Mr. Smith meets the IP criteria.
- (DENOM = YES) Mr. Smith meets the DENOM criteria.
- (DENEX = NO) By the positive evidence principle, Mr. Smith does not meet the DENEX criteria.
- (NUMER = NO) By the positive evidence principle, Mr. Smith does not meet the NUMER criteria.
- (NUMEX = NO)
- (DENEXCEP = NO) By the positive evidence principle, Mr. Smith does not meet the DENEXCEP criteria.

Mr. Johnson is 85 years old and has an active diagnosis of diabetes. There is no mention of blindness in his chart. He has a documented dilated eye exam for diabetic retinopathy and is confined to bed in a community where mobile eye exam imaging is unavailable.

- (IP = YES) Mr. Johnson meets the IP criteria.
- (DENOM = YES) Mr. Johnson meets the DENOM criteria.
- (DENEX = NO) By the positive evidence principle, Mr. Johnson does not meet the DENEX criteria.
- (NUMER = YES) Mr. Johnson meets the NUMER criteria.
- (NUMEX = NO)
- (DENEXCEP = NO) By definition, Mr. Johnson does not meet the DENEXCEP criteria because DENEXCEP criteria are not applicable to those meeting the NUMER criteria.

Example #2: Inverse Proportion Measure

A fictitious inverse proportion measure (i.e., where improvement is a decrease in the rate) defines the population criteria.

- IP: patients aged 8 to 65 who gave birth
- DENOM: equals IP
- DENEX: patients giving birth to an infant with gestational age < 37 weeks
- NUMER: patients with medical induction of labor or C-section
- NUMEX: patients in active labor or with spontaneous rupture of membranes before induction of labor or C-section
- DENEXCEP: none

Mrs. Jones is a 31-year-old woman who gave birth at 37 weeks' gestation. She has a medical induction of labor with no prior evidence of active labor or spontaneous rupture of membranes.

- (IP = YES) Mrs. Jones meets the IP criteria.
- (DENOM = YES) Mrs. Jones meets the DENOM criteria.
- (DENEX = NO) Mrs. Jones does not meet the DENEX criteria.
- (NUMER = YES) Mrs. Jones meets the NUMER criteria.
- (NUMEX = NO) By the positive evidence principle, Mrs. Jones does not meet the NUMEX criteria.
- (DENEXCEP = NO)

Mrs. Thompson is 31 years old and had a C-section at 38 weeks after a spontaneous rupture of membranes.

- (IP = YES) Mrs. Thompson meets the IP criteria.
- (DENOM = YES) Mrs. Thompson meets the DENOM criteria.
- (DENEX = NO) Mrs. Thompson does not meet the DENEX criteria.
- (NUMER = YES) Mrs. Thompson meets the NUMER criteria.
- (NUMEX = YES) Mrs. Thompson meets the NUMEX criteria.
- (DENEXCEP = NO)

Mrs. Hill is 31 years old and gave birth at 36 weeks with induction after a spontaneous rupture of membranes.

- (IP = YES) Mrs. Hill meets the IP criteria.
- (DENOM = YES) Mrs. Hill meets the DENOM criteria.
- (DENEX = YES) Mrs. Hill meets the DENEX criteria.
- (NUMER = NO) By definition, Mrs. Hill does not meet the NUMER criteria because NUMER criteria are not applicable to those meeting the DENEX criteria.
- (NUMEX = NO) By definition, Mrs. Hill does not meet the NUMEX criteria because NUMEX criteria are only applicable to those meeting the NUMER criteria.
- (DENEXCEP = NO)

Example #3: Measure Aggregate Calculations

Aggregate scores⁽ⁱ⁾ are simply the counts of individuals in each population. Thus, the aggregate IP is the count of individuals meeting the IP criteria.

Building on Example #1, by counting all individuals within the population, the aggregate counts are determined.

- Target/Initial population: N=150 (e.g., 150 patients meet the IP criteria)
- Denominator: N=150

- Denominator exclusion: N=20 (meet DENOM and meet DENEX)
- Numerator: N=75 (meet DENOM, not in DENEX, and meet NUMER criteria)
- NUMEX: N=0
- Denominator exception: N=5 (meet DENOM, not in DENEX, not in NUMER, and meet the DENEXCEP criteria)
- Performance rate = $(\text{NUMER} - \text{NUMEX}) / (\text{DENOM} - \text{DENEX} - \text{DENEXCEP}) = (75-0) / (150-20-5) = 0.6$

2 CONTINUOUS VARIABLE CALCULATIONS

This section provides further guidance on the precise mathematical relationships between populations in a CV measure and the process for use in determining individual and aggregate scores.

In a CV measure, each individual value for the measure can fall anywhere along a continuous scale, for example, mean number of minutes between presentation of chest pain to the time of administration of thrombolytic medications.

2.1 ELEMENTS OF CONTINUOUS VARIABLE MEASURES

There is a fixed mathematical relationship between the populations in a CV measure, as shown in [Figure 3](#):




Figure 3. CV Measure Populations





Figure 4. Population Criteria for Continuous Variable Measures (HL7, n.d.)

Figure 4 illustrates the flow from target/initial population to measure observation. Note the figure references electronic clinical quality measures (eQMs) ^①, but is valid for other measure types.

We define the CV measure query process as follows:

- IP: Identify those cases that meet the IP criteria.
- Measure population (MSRPOPL): Identify the subset of the IP that meet the MSRPOPL criteria.
- Measure population exclusion (MSRPOPLEX): Identify the subset of the MSRPOPL that meet the MSRPOPLEX criteria.

For a CV measure, an individual observation is determined for each case in the MSRPOPL and not in the MSRPOPLEX.


 eQIM Clinical Quality Language (CQL) ¹ logic ¹ should not explicitly remove the MSRPOPLEX cases because the execution environment for the eQIM is responsible for excluding the MSRPOPLEX cases and calling the Measure Observation function for each of the remaining cases.

Per Requirement 13 of the [HL7® Version 3 Implementation Guide: CQL-based Health Quality Measure Format \(HQMF\) Release 1, Standard for Trial Use \(STU\) 3 – US Realm](#) ¹, aggregation rules are mandatory: “methodCode shall be populated to indicate the aggregation method for the measure.”

2.2 CONTINUOUS VARIABLE AGGREGATE CALCULATIONS

Aggregate scores for CV measures are more complex than for proportion measures in that they are more than just the counts of cases in each population. In addition to the identification of measure population(s), CV measures define observations made on cases falling into various populations. The measure developer then aggregates the individual observations, after removing the exclusions, according to aggregation rules specific to each measure.

In CV measures, the measure developer should make an individual observation for each case falling into the measure population and not in the measure population exclusion. Then the measure developer uses the remaining individual observations to calculate the aggregate CV, which constitutes the measure “score.”

 It is important to note that Quality Reporting Document Architecture (QRDA) ¹ Category I can send observations on the individual, whereas QRDA Category III can send aggregate calculations.

2.3 CV MEASURE EXAMPLE

A fictitious CV measure defines the population criteria:

- IP: All inpatient encounters ¹ ending during the measurement period with length of stay fewer than or equal to 120 days
- MSRPOPL: Inpatient encounters preceded by an emergency department (ED) visit
- MSRPOPLEX: none
- Individual observations: Time in minutes from ED admission to ED discharge for patients admitted to the facility from the ED
- MSRPOPL observation: Number of minutes in the ED

Aggregation rules:

 For eQIMs, HQMF ¹ methodCode specifies median.

Mr. Jones is a 75-year-old man admitted to the hospital from the ED. He spent 90 minutes in the ED.

- (IP = YES) Mr. Jones has inpatient encounters that meet the IP criteria.
- (MSRPOPL = YES) Mr. Jones meets the MSRPOPL criteria.
- (MSRPOPLEX = NO)
- Individual observations for Mr. Jones: 90 minutes
- MSRPOPL observation: 90 minutes

Mr. James is 75 years old, admitted directly to the hospital from an outside facility.

- (IP = YES) Mr. James has inpatient encounters that meet the IP criteria.
- (MSRPOPL = NO) By the positive evidence principle, Mr. James does not meet the MSRPOPL criteria.
- (MSRPOPLEX = NO)
- Individual observations for Mr. James: none
- MSRPOPL observation: N/A (Mr. James has no inpatient encounters that meet the MSRPOPL criteria).

Measure aggregate calculations:

Building upon the previous examples, but now considering all encounters within the population, the aggregate counts are determined.

- IP: N = 150 (i.e., 150 encounters meet the IP criteria)
- MSRPOPL: N = 120
- MSRPOPLEX: N = 0

CV Score: 96 minutes (median of all individual Measure Population observations of time spent in the ED).

3 RATIO MEASURE CALCULATIONS

This section provides further guidance on the precise mathematical relationships between populations in a ratio measure. The measure developer can use this process to determine individual and aggregate scores.

The measure developer can derive a ratio measure by dividing a count of one type of data by a count of another type of data (e.g., the number of patients with central lines who develop infection divided by the number of central line days). A ratio measure differs from a proportion measure because in a ratio measure the numerator is not a subset of the denominator population. Although the numerator and denominator populations may be related or may overlap, these populations do not have a superset/subset relationship.

3.1 ELEMENTS OF A RATIO MEASURE

There is a fixed mathematical relationship between the populations in a ratio measure, as shown in

[Figure 5](#). Ratio measures are often two CV calculations for related populations (e.g., median ED waiting time for the index hospital; median ED waiting time for the region in which the hospital is located as the numerator).




Figure 5. Ratio Measure Populations




Figure 6. Population Criteria for Ratio Measures - Numerator (HL7, n.d.)

Figure 6 illustrates the flow from target/initial population to the numerator and numerator exclusion. Figure 7 illustrates the flow from target/initial population to the denominator and denominator exclusion.




Figure 7. Population Criteria for Ratio Measures - Denominator (HL7, n.d.)

From these relationships and definitions, the measure developer defines the ratio measure query process:

- IP: Identify those cases that meet the IP criteria. (Some ratio measures will require multiple IPs: one for the numerator and one for the denominator).
- DENOM: Identify the subset of the IP that meet the DENOM criteria.
- DENEX: Identify the subset of the DENOM that meet the DENEX criteria.
- NUMER: Identify the subset of the IP that meet the NUMER criteria.
- NUMEX: Identify the subset of the NUMER that meet the NUMEX criteria.

The measure developer then computes individual observations for ratio measures both for the denominator and numerator populations:

- For each case in the DENOM and not in the DENEX, determine the individual DENOM observations.

- For each case in the NUMER and not in the NUMEX, determine the individual NUMER observations.

3.2 RATIO MEASURE AGGREGATE CALCULATIONS

As with CV measures, aggregate scores for ratio measures are more than just the counts of cases in each population. In addition to the identification of measure population(s), the measure developer should make ratio measures define observations on cases falling into various populations and then aggregate these individual observations according to aggregation rules specific to each measure.

In ratio measures, for each population, the measure developer should use individual observations, e.g., measurements or calculations, for denominator and numerator cases, and then use them to calculate the aggregate ratio:

$$\text{Ratio} = \text{aggregate NUMER} / \text{aggregate DENOM}$$

Calculate the aggregate DENOM using individual observations for all cases in the DENOM and not in the DENEX, and calculate the aggregate NUMER using individual observations for all cases in the NUMER and not in the NUMEX.

3.3 RATIO MEASURE EXAMPLE

A fictitious ratio measure looking at the rate of central line blood stream infections (i.e., the ratio of number of central line infections per total central line days) defines the population criteria as follows:

- IP: All hospitalized patients with a laboratory confirmed bloodstream infection
- DENOM: Presence of central line or umbilical catheter for >2 calendar days
- DENEX: Patient is immunosuppressed.
- NUMER: Central line blood stream infection
- NUMEX: Deem infection to be a contaminant

Individual Observations:

- DENOM observation: Number of hospital days with central line
- NUMER observation: Fully define the observation in this case by the population criteria (i.e., the observation is simply whether the case met the population criteria or not).

Aggregation Rules:

- DENOM aggregation: Sum of number of hospital days with central line – DENEX
- NUMER aggregation: Count of NUMER – NUMEX

Mr. Jones is 75 years old and was in the hospital for 7 days. He had a central line in place for 5 days. There is no mention of immunosuppression in his chart. There is no mention of a central line blood stream infection in his chart.

- (IP = YES) Mr. Jones meets the IP criteria.
- (DENOM = YES) Mr. Jones meets the DENOM criteria.
- (DENEX = NO) By the positive evidence principle, Mr. Jones does not meet the DENEX criteria.
- (NUMER = NO) Mr. Jones does not meet the NUMER criteria.

- (NUMEX = NO) By definition, Mr. Jones does not meet the NUMEX criteria because NUMEX criteria are only applicable to those meeting the NUMER criteria.

Individual observations for Mr. Jones:

- DENOM observation: 5
- NUMER observation: N/A (Mr. Jones does not meet the NUMER criteria)

Mr. James is 75 years old and was in the hospital for 24 days. He had a central line in place for 17 days. There is no mention of immunosuppression in his chart. He has a documented central line blood stream infection while the central line was in place. There is no mention that the infection is a contaminant.

- (IP = YES) Mr. James meets the IP criteria.
- (DENOM = YES) Mr. James meets the DENOM criteria.
- (DENEX = NO) By the positive evidence principle, Mr. James does not meet the DENEX criteria.
- (NUMER = YES) Mr. James meets the NUMER criteria.
- (NUMEX = NO) By the positive evidence principle, Mr. James does not meet the NUMEX criteria.

Individual observations for Mr. James:

- DENOM observation: 17
- NUMER observation: Mr. James meets the NUMER criteria.

Measure aggregate calculations


Building from the examples above, by considering all cases within the population, the aggregate counts are determined.

- IP: N = 150 (i.e., 150 cases meet the IP criteria)
- DENOM: N = 20 of the 150 had laboratory confirmed blood stream infection with central lines.
- DENEX: N = 2
- NUMER: N = 6 (Mr. James was one of the six numerator cases.)
- NUMEX: N = 1

Aggregate DENOM: 108 (In this example, 108 is an assumed sum of central line days across all cases in the DENOM and not in the DENEX).


Aggregate NUMER: 5 (total number of central line blood stream infections, excluding those deemed to be contaminants).

Ratio = aggregate NUMER / aggregate DENOM = 5 / 108 = 0.046

 Find additional information about eQMs and proportion, ratio, and CV measure calculation at the [Electronic Clinical Quality Improvement \(eCQI\) Resource Center](#).

4 KEY POINTS

The measure calculation is an important part of measure specification^①: it defines the mathematical relationships among measure components (numerator, denominator, denominator and numerator exclusions, and denominator exceptions) and dictates how the measure score should be interpreted. Although there are many ways to calculate a measure score, the most common are proportions, CVs, and ratios. Proportion measures are those in which the numerator is a subset of the denominator. They

differ from ratio measures because, although numerator and denominator populations may be related for ratio measures, they do not have a superset/subset relationship. CV measures, on the other hand, allow for individual scores that can fall anywhere along a continuous scale (e.g., mean number of minutes between arrival and admission). Population flow diagrams easily depict each of these calculation methods or calculation algorithm  that illustrates the relationships among measure components to ensure the calculation of measure scores is done in a standardized way.

REFERENCES

Electronic Clinical Quality Improvement Resource Center. (n.d.). *eCQI resource center*. Retrieved March 22, 2021, from <https://ecqi.healthit.gov/>

HL7 International. (n.d.). *Quality measure implementation guide*. Retrieved April 5, 2021, from <http://build.fhir.org/ig/HL7/cqf-measures/measure-conformance.html#ratio-measures>

HL7 International. (2018). *Health Level Seven International (HL7) version 3 implementation guide: CQL-based health quality measure format (HQMF) release 1, standard for trial use (STU) 3 - US realm*. Retrieved March 22, 2021, from <http://www.hl7.org/dstucomments/showdetail.cfm?dstuid=230>